



20 DISTRIBUTION/AVAILABILITY OF ABSTRACT

UNCLASSIFIED

21 ABSTRACT SECURITY CLASSIFICATION

UNCLASSIFIED

226 NAME OF RESPONSIBLE INDIVIDUAL

Dr. William O. Berry

21 ABSTRACT SECURITY CLASSIFICATION

UNCLASSIFIED

226 OFFICE SYMBOL (Include Area Code: (202) 767-5021

NL

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE

UNCLASSIFIED

Routtenberg

Annual Technical Report

AF 0SR 87-0042

Annual Technical/Scientific Report

- 1. <u>Project Period</u> The project period includes October 1, 1986 to September 30, 1987. The present report is being filed in August, 1987.
 - 2. Summary

The regulation of synaptic reactivity by protein kinase C and its substrate proteins has been studied using the long-term potentiation paradigm (LTP). In the past year we have studied the effects of protein kinase C activators and inhibitors on durability of synaptic reactivity. The main conclusion to be drawn is that protein kinase C is necessary but not sufficient for the enhanced durability. In combination with a neural signal, however, PKC demonstrates a profound synergism. Synergism is also observed in the analysis of metal ion regulation of protein kinase C activity. Calcium and zinc interact in their effect on the enzyme in a



3. Statement of Work

bidirectional manner (see below).

CONTINUE CONTINUE CONTINUE SOCION DISCOUNT INSTANCE AND SOUD SESSION SESSION CONTINUE

The research objectives during this period were to:

- a. Study effects of protein kinase C inhibitors
- b. Study activators of PKC in the synaptic zone
- c. Study metal ion regulation of PKC
- 4. Status of Research
 Significant accomplishments made during this period were:
- a. Effect of inhibitors

We used three separate inhibitors of protein kinase C activity,

-1-



こうじょうじょう またくどうしょうじょ またなななななな しゅうしきしきしき

polymyxin B, mellitin and H-7, each with a different mechanism of action. Application was made by micro-pressure ejection into the molecular layer of the dentate gyrus before or after LTP. The major result of this study was that inhibitors had no effect on the initiation of LTP but completely eliminated the enhanced response 10-15 min after its initiation. This provides strong support for our view that PKC plays a critical role in the maintenance but not the initiation of LTP.

b. Study of PKC activators (PDBu and oleate)

A crucial question in the analysis of the role of PKC in synaptic reactivity is the site of action of the compound. Indirect evidence suggested a synaptic site since PKC is found in high concentration there. To assess this view directly we compared application dosages required to facilitate synaptic reactivity duration in the dentate hilus, a nearby site, and the molecular layer of the dentate, 100 micra from the granule cells, precisely the point where perforant path terminals synapse. We have found that only 10-16% of the dosage is required when the application, iontophoretic or micro-pressure, is made at the synaptic zone. This provides strong support for the synaptic site of action of these protein kinase C activators.

c. Metal ion regulation of protein kinase C activity

Recent evidence describing the primary structure of protein kinase C by several laboratories indicates several different motifs: ATP-binding, Ca-binding, kinase domain, zinc "fingers". This suggested the possibility that both zinc as well as calcium might regulate protein kinase C activity.

Since we have recently discovered that protein kinase C can be activated in the absence of calcium it was now feasible to sstudy the effects of zinc both in the presence and the absence of calcium. A novel mechanism for regulating protein kinase C activity was discovered in which zinc ions, found in highest concentration in the hippocampus, enhance protein kinase C activity at low calcium levels. At higher levels of calcium, zinc inhibits. We propose a model of protein kinase C with a low and high calcium affinity binding sites and a distinct zinc binding site.

- 5. Articles published, accepted for publication and submitted.
- Routtenberg, A., Colley, P., Linden, D., Lovinger, D., and Murakami, K.

 Phorbol ester promotes growth of synaptic plasticity. Brain Research,

 1986, 378, 374-378.
- Linden, D., Murakami, K., and Routtenberg, A. A newly discovered protein kinase C activator (oleic acid) preserves synaptic plasticity and promotes the growth of long-term potentiation. Brain Research, 1986, 379, 358-363.
- Gispen, W.H. and Routtenberg, A. (Eds.). <u>Phosphoproteins in the Nervous</u>

 System. Elsevier/North Holland, Amsterdam, 1986, pg. 211-234.
- Lovinger, D., Akers, R., Colley, P., Linden, D., and Routtenberg, A.

 Direct relation of synaptic plasticity to phosphorylation of membrane protein F1: Potential association with protein kinase C translocation.

 Brain Research, 1986, 399, 205-211.
- Routtenberg, A. A new theory of Alzheimer's disease. Arts and Sciences, Northwestern University, 1986, 9, 2-6.

- Chan, S.Y., Murakami, K., and Routtenberg, A. Phosphoprotein F1:

 Purification and characterization of a brain kinase C substrate related to plasticity. J. Neuroscience, 1986, 6, 3618-3627.
- Murakami, K., Chan, S.Y., and Routtenberg, A. Protein kinase C activation by cis-fatty acid in the absence of Ca2+ and phospholipids. <u>J. biol.</u> Chem., 1986, 261, 15424-15429.
- Collier, T.J., Quirk, G.S., and Routtenberg, A. Separable roles of hippocampal granule cells in forgetting and pyramidal cells in remembering spatial information. Brain Research, 1987, in press.
- Lovinger, D.M. and Routtenberg, A. Protein F1 and protein kinase C may regulate the persistence of synaptic potentiation in the hippocampus. In: Ehrlich, Y.H., Berry, W., and Lennox, R. (Eds.). Molecular Mechanisms of Neuronal Responsitivity. To appear in: Advances in Experimental Biology and Medicine, New York: Plenum, 1987, in press.
- Ruth, R.E., Collier, T.J., and Routtenberg, A. Topography between the entorhinal cortex and the dentate septotemporal axis in rats:

 II. Lateral entorhinal projecting cells. <u>J. comp. Neurol.</u>, 1987, in press.
- Routtenberg, A. (1987) Phospholipid and fatty acid regulation of signal transduction at synapses: Potential role for protein kinase C in information storage. To appear in: R.J. Wurtman (Ed.). "Alzheimer's Disease: Advances in Basic Research and Therapies". pp. 1-10.
- Nelson, R.B., Friedman, D.P., O'Neill, J.B., Mishkin, M., and Routtenberg,
 A. Gradients of PKC substrate phosphorylation along the primate visual

- cortical processing pathway: Highest levels are in visual memory areas. Brain Research, 1987, in press.
- Akers, R.F. and Routtenberg, A. Calcium-promoted translocation of protein kinase C to synaptic membranes: relation to the phosphorylation of an endogenous substrate (Protein F1) involved in synaptic plasticity. J. Neurosci., 1987, in press.
- Snipes, G.J., Chan, S., McGuire, C.B., Costello, B.R., Routtenberg, A., Norden, J.J., and Freeman, J.A. Evidence that GAP-43, a growth-related protein, and Fl, a synaptic plasticity-associated protein, are identical. J. Neurosci., 1987, in press.
- Linden, D., Shew, F.-S., Murakami, K., and Routtenberg, A. Cis fatty acid regulation of synaptic potentiation: Relation to phospholipase A2 and protein kinase C activation. J. Neurosci., 1987, in press.
- Murakami, K., Whiteley, M.K., and Routtenberg, A. Regulation of protein kinase C activity by cooperative interaction of Zn2+ and Ca2+. J. Biol. Chem., 1987, in press.
- Benowitz, L.I. and Routtenberg, A. A membrane phosphoprotein associated with neural development, axonal regeneration, phospholipid metabolism, and synaptic plasticity. Submitted.
- Lovinger, D.M. and Routtenberg, A. Synapse specific protein kinase C activation enhances maintenance of long-term potentiation in rat hippocampus. Submitted.
- Nelson, R.B., Linden, D.J., and Routtenberg, A. Two protein kinase C substrates and two vesicle-associated phosphoproteins are directly

THE EXCEPTION OF THE PRODUCT OF THE

- correlated with persistence of long-term potentiation: A quantiative analysis of two-dimensional gels. Submitted.
- Nelson, R.B., Hyman, C., Pfenninger, K.H., and Routtenberg, A. Two protein kinase C substrates directly correlated with persistence of long-term potentiation in adult rat brain are the major phosphoproteins found in nerve growth cones. Submitted.
- Chan, S.Y., Nelson, R.B., Murakami, K., and Routtenberg, A. Protein kinase C substrate (F1) phosphorylation: Phospholipid-independent, CA2+-enhanced CIS-fatty acid activation. Submitted.
- Barnes, C.A., Mizumori, S.J.Y., Lovinger, D.M., Sheu, F.-S., Murakami, K., Chan, S.Y., Linden, D.J., Nelson, R.B., and Routtenberg, A. Selective decline in protein F1 phosphorylation in hippocampus of senescent rats. Submitted.
- Nelson, R.B. and Routtenberg, A. Contrasting roles of a brain specific protein kinase C substrate: Has protein F1 evolved a new function in CNS of higher vertebrates? In: Proceedings of the IUPS Satellite
 Plasticity (C. Woody, Ed.), Seattle, WA. July 9-12, 1986. In press.
- Rosenthal, A., Chan, S.Y., Henzel, W., Haskell, C., Kuang, W.-J., Chen, E., Wilcox, J.N., Ullrich, A., Goeddel, D.V., and Routtenberg, A. Primary structure and mRNA localization of protein Fl, a growth-related protein kinase C substrate, associated with synaptic plasticity. Submitted.
- Nelson, R. and Routtenberg, A. The protein F1/protein kinase C module and neurite growth: Potential utility in facilitating brain

Routtenberg

transplantation. In: <u>Progress in Brain Research</u> (J. Slarek and D. Gash, Eds.), Vol. 82, 1988, in press.

Personnel

<u>Name</u>	<u>Title</u>	Dates of Service	% Effort
A. Routtenberg	Professor/PI	9/83-present	25%
S. Chan	Res. Neurobiologist	2/84-present	25%
K. Murakami	Res. Neurobiologist	4/84-present**	25%
P. Colley	Grad. Res. Asst.	7/83-present	50%
D. Linden	Grad. Res. Asst.	9/84-present	50%
D. Lovinger	Grad. Res. Asst.	7/83-present*	50%
R. Nelson	Grad. Res. Asst.	7/83-present*	50%
F. Sheu	Grad. Res. Asst.	9/85-present	50%
K. Wong	Grad. Res. Asst.	4/86-present	50%

^{* -} Ph.D. awarded 6/87

- ** Assistant Professor, University of Buffalo, September, 1987.
 - 7. Coupling Activities
- Colley, P. and Routtenberg, A. Hypothesis: Protein kinase C activation acts synergistically with a calcium-mediated event to induce long-lasting synaptic changes in the hippocampus. Soc. Neurosci., 1986, 12, 1168.
- Lovinger, D., Barnes, C.A., Mizumori, S.J.Y., Chan, S.Y., Linden, D.,

 Murakami, K., Sheu, F.-S., and Routtenberg, A. Protein F1, previously
 related to synaptic plasticity, exhibits decreased phosphorylation in

- senescent rat hippocampus. Soc. Neurosci. 1986, 12, 1168.
- Nelson, R.B., Friedman, D.P., Mishkin, M., and Routtenberg, A. Protein kinase C substrate phosphorylation in primate cerebral cortex (E.G. Protein F1) is increased in those stages of the occipitotemporal visual processing pathway important for information storage. Soc. Neurosci., 1986, 12, 1168.
- Chan, S.Y., Murakami, KI., and Routtenberg, A. Characterization of purified protein Fi. a specific protein kinase C substrate related to synaptic plasticity. <u>Soc. Neurosci.</u>, 1986, 12, 1168.
- Linden, D.J., Murakami, K., and Routtenberg, A. Oleic acid, a protein kinase C activator, enhances hippocampal long-term potentiation. <u>Soc.</u>
 Neurosci., 1986, 12, 1169.
- Murakami, K., Chan, S.Y., and Routtenberg, A. Protein kinase C (PKC) activation by cis-fatty acid is distinct from previously described phospholipid and Ca2+ activation. Soc. Neurosci., 1986, 12, 1169.
- Snipes, G.J., Freeman, J.A., Costello, B., Chan, S., and Routtenberg, A. A growth associated protein, GAP-43, is immunologically and structurally related to the plasticity associated protein, protein Fl. Soc.

 Neurosci., 1986, 422.
- Routtenberg, A. Phospholipid and fatty acid regulation of signal transduction at synapses: Potential role for protein kinase C in information storage. Invited Lecture: Fourth meeting of the International Study Group on The Pharmacology of Memory Disorders Associated with Aging. Hotel Zurich, Zurich, Switzerland, January

16-18, 1987.

- Colley, P. and Routtenberg, A. Induction, enhancement and blockade of hippocampal synaptic potentiation: Dose-dependent effects of phorbol 12,13-dibutyrate (PDBu). Soc. Neurosci., 1986, 13, in press.
- Lovinger, D., Wong, K., Murakami, K., and Routtenberg, A. Protein kinase C (PKC) inhibitors prevent maintenance of hippocampal long-term potentiation (LTP). <u>Soc. Neurosci.</u>, 1986. <u>13</u>, in press.

END FB. 198 1